




# Genome Sequence of the Mycotoxigenic Crop Pathogen *Fusarium proliferatum* Strain ITEM 2341 from Date Palm

Bandar F. Almiman,<sup>a</sup> Taiwo A. Shittu,<sup>a</sup> Sreenivasaprasad Muthumeenakshi,<sup>a</sup>  Riccardo Baroncelli,<sup>a,b</sup> Surapareddy Sreenivasaprasad<sup>a</sup>

<sup>a</sup>School of Life Sciences, Institute of Biomedical and Environmental Science and Technology (iBEST), University of Bedfordshire, Luton, United Kingdom

<sup>b</sup>University of Salamanca, Salamanca, Spain

**ABSTRACT** *Fusarium proliferatum* is a widely distributed fungal pathogen associated with more than 26 crop species important in global food security. Its strong mycotoxigenic capability with potential impacts on human and animal health is well recognized. In this work, we report the draft genome sequence of *F. proliferatum* strain ITEM 2341, originally isolated from date palm, providing a platform for further comparative and functional genomic investigations.

*Fusarium proliferatum* is an emerging fungal pathogen belonging to the *F. fujikuroi* species complex within the *Fusarium* genus. The *F. fujikuroi* complex contains an estimated 50 species (1). *F. proliferatum* is widely distributed and associated with economically important crops such as rice, maize, sugarcane, date palm, onion, and tomato (2). *F. proliferatum* is able to cause different types of diseases, such as dieback, wilt, and rot, at different stages of the crop affecting various parts. *F. proliferatum* can also colonize diverse types of host plants, including maize and orchids, without displaying visible symptoms (3, 4). In addition to its direct impact on crop production, *F. proliferatum* is well recognized to produce high levels of mycotoxins, such as fumonisins, which are secondary metabolites known to be carcinogenic (4, 5). *F. proliferatum* ITEM 2341 was originally isolated in 1998 from the roots of date palm (*Phoenix dactylifera*) in Buraydah (Al-Qassim, Saudi Arabia; <http://server.ispa.cnr.it/ITEM/Collection/>).

The *F. proliferatum* strain ITEM 2341 was grown in potato dextrose broth for 5 days at 20°C. The mycelium was harvested using sterile muslin cheesecloth, air dried at room temperature for 15 min, and ground into a fine powder in liquid nitrogen with a sterile pestle and mortar. Genomic DNA was extracted from the mycelial powder using the GenElute plant genomic DNA miniprep kit (product number G2N350-1KT; Sigma, UK). The genome of *F. proliferatum* ITEM 2341 was sequenced using a combination of paired-end and mate pair libraries with the Illumina MiSeq platform using the service provider at the University of Cambridge, United Kingdom. Paired-end and mate pair libraries were prepared using the Illumina TruSeq PCR-free kit and the Illumina Nextera mate pair kit, respectively. The average length of DNA fragments used was 550 bp for paired-end sequencing and 2.5 to 4 kb for mate pair sequencing. The numbers of reads generated from the paired-end and mate pair libraries were 6,182,938 and 11,135,290, respectively. At 300 bases per read length, the total number of bases generated was ~5.08 billion. FastQC version 0.11.5 was used to check the quality of the reads, and low-quality bases with a Phred score of less than Q20 as well as adaptor sequences were filtered using the BBDuk plugin within Geneious version 9.1.5. The curated reads were assembled using SPAdes version 3.5.0 (6), providing an approximate coverage of 111×.

The nuclear genome of *F. proliferatum* strain ITEM 2341 was assembled in 104 scaffolds with a total assembly size of 45.50 Mb (48.29% GC content). The genome

Received 9 July 2018 Accepted 9 August 2018 Published 6 September 2018

**Citation** Almiman BF, Shittu TA, Muthumeenakshi S, Baroncelli R, Sreenivasaprasad S. 2018. Genome sequence of the mycotoxigenic crop pathogen *Fusarium proliferatum* strain ITEM 2341 from date palm. Microbiol Resour Announc 7:e00964-18. <https://doi.org/10.1128/MRA.00964-18>.

**Editor** Antonis Rokas, Vanderbilt University

**Copyright** © 2018 Almiman et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Surapareddy Sreenivasaprasad, [ss.prasad@beds.ac.uk](mailto:ss.prasad@beds.ac.uk).

assembly included the following parameters: an  $N_{50}$  value of 1,822,579 bp, an  $L_{50}$  value of 9, and a maximum scaffold size of 3,489,699 bases. Genome assembly quality assessment using BUSCO version 3.0.2 (7) with the Sordariomyceta data set (3,725 genes) showed the presence of 3,691 genes (99.1% gene content). The nuclear genome was annotated using the MAKER2 pipeline (8), and 14,379 protein-coding gene models were predicted. SignalP version 4.1 (9) analysis revealed 1,488 secreted proteins (10.35% of the predicted proteome). All software used for the assembly and analysis of the genome was set at default parameters.

The genome sequence of *F. proliferatum* ITEM 2341 represents a useful platform for further comparative genomic analysis of the adaptive divergence in this emerging pathogen (10–12).

**Data availability.** The whole-genome shotgun project has been deposited in DDBJ/ENA/GenBank under the accession number PKMI00000000 (BioSample number SAMN08122925; BioProject number PRJNA420865; NCBI Sequence Read Archive [SRA] accession numbers SRX4488677 and SRX4488678 [raw reads]) and released for access by the research community. The version described in this paper is PKMI01000000.

## ACKNOWLEDGMENTS

Funding support to Bandar F. Almiman's Ph.D. program by the Saudi Arabian government through their Cultural Bureau in London, United Kingdom, is gratefully acknowledged.

We thank Giuseppe Cozzi and Giancarlo Perrone of the Agri-Food Microbial Culture Collection (ITEM), Institute of Sciences of Food Production, National Research Council, Italy, for providing the *F. proliferatum* isolate ITEM 2341. We gratefully acknowledge the bioinformatics resources support from the research group led by Mike Thon, University of Salamanca, Spain.

## REFERENCES

- O'Donnell K, Ward TJ, Robert VARG, Crous PW, Geiser DM, Kang S. 2015. DNA sequence-based identification of *Fusarium*: current status and future directions. *Phytoparasitica* 43:583–595. <https://doi.org/10.1007/s12600-015-0484-z>.
- Proctor RH, Desjardins AE, Moretti A. 2010. Biological and chemical complexity of *Fusarium proliferatum*. In Strange RN, Gullino ML (ed), *The role of plant pathology in food safety and food security in the 21st century*. Springer, Dordrecht, The Netherlands.
- Munkvold GP, Hellmich RL, Showers WB. 1997. Reduced *Fusarium* ear rot and symptomless infection in kernels of maize genetically engineered for European corn borer resistance. *Phytopathology* 87:1071–1077. <https://doi.org/10.1094/PHYTO.1997.87.10.1071>.
- Tsavkelova EA, Bömké C, Netrusov AI, Weiner J, Tudzynski B. 2008. Production of gibberellic acids by an orchid-associated *Fusarium proliferatum* strain. *Fungal Genet Biol* 45:1393–1403. <https://doi.org/10.1016/j.fgb.2008.07.011>.
- Scott PM. 2012. Recent research on fumonisins: a review. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 29:242–248. <https://doi.org/10.1080/19440049.2010.546000>.
- Bankovich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31:3210–3212. <https://doi.org/10.1093/bioinformatics/btv351>.
- Holt C, Yandell M. 2011. MAKER2: an annotation pipeline and genome-database management tool for second-generation genome projects. *BMC Bioinformatics* 12:491. <https://doi.org/10.1186/1471-2105-12-491>.
- Petersen TN, Brunak S, von Heijne G, Nielsen H. 2011. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods* 8: 785–786. <https://doi.org/10.1038/nmeth.1701>.
- Ma LJ, van der Does HC, Borkovich KA, Coleman JJ, Daboussi MJ, Di Pietro A, Dufresne M, Freitag M, Grabherr M, Henrissat B, Houterman PM, Kang S, Shim WB, Woloshuk C, Xie X, Xu JR, Antoniw J, Baker SE, Bluhm BH, Breakspear A, Brown DW, Butchko RA, Chapman S, Coulson R, Coutinho PM, Danchin EG, Diener A, Gale LR, Gardiner DM, Goff S, Hammond-Kosack KE, Hilburn K, Hua-Van A, Jonkers W, Kazan K, Kodira CD, Koehrsen M, Kumar L, Lee YH, Li L, Manners JM, Miranda-Saavedra D, Mukherjee M, Park G, Park J, Park SY, Proctor RH, Regev A, Ruiz-Roldan MC, Sain D, et al. 2010. Comparative genomics reveals mobile pathogenicity chromosomes in *Fusarium*. *Nature* 464:367–373. <https://doi.org/10.1038/nature08850>.
- Zapparata A, Da Lio D, Somma S, Vicente Muñoz I, Malfatti L, Vannacci G, Moretti A, Baroncelli R, Sarrocco S. 2017. Genome sequence of *Fusarium graminearum* ITEM 124 (ATCC 56091), a mycotoxigenic plant pathogen. *Genome Announc* 5:e01209-17. <https://doi.org/10.1128/genomeA.01209-17>.
- Niehaus EM, Münsterkötter M, Proctor RH, Brown DW, Sharon A, Idan Y, Oren-Young L, Sieber CM, Novák O, Pěnčík A, Tarkowská D, Hromádová K, Freeman S, Maymon M, Elazar M, Youssef SA, El-Shabrawy ES, Shalaby AB, Houterman P, Brock NL, Burkhardt I, Tsavkelova EA, Dickschat JS, Galuszka P, Güldener U, Tudzynski B. 2016. Comparative “omics” of the *Fusarium fujikuroi* species complex highlights differences in genetic potential and metabolite synthesis. *Genome Biol Evol* 8:3574–3599. <https://doi.org/10.1093/gbe/evw259>.